

## **OBSTETRICS/GYNECOLOGY**

### **⌘ A Randomized Trial of Thyroxine Therapy for Subclinical Hypothyroxinemia Diagnosed in Pregnancy**

Developing babies need thyroid hormones while they are in the womb in order to grow and develop normally. Women with too much (called hyperthyroidism) or too little (called hypothyroidism) thyroid hormone often have problems in pregnancy that can be prevented or reduced by treatment. You do not have either of these conditions.

This study is being done to find out whether every pregnant woman should be tested for mildly under-active thyroid gland and then be treated if she does. Some women have mildly under-active thyroid glands, a condition where they are normal and feel fine, but have thyroid blood tests that are not quite right.

This clinical trial is being conducted at: Mount Carmel St. Ann's

For more information about this clinical trial call: 614-898-5587 or 614-293-4652

### **⌘ A 24-Week, Randomized, Double-Blind, Placebo Controlled, Safety and Efficacy Trial of Flibanserin 50 and 100 Milligrams Each Evening in Premenopausal Women with Hypoactive Sexual Desire Disorder**

This clinical trial is being conducted to find out if an investigational drug called "flibanserin" is safe and effective for the treatment of premenopausal women with Hypoactive Sexual Desire Disorder or HSDD. Flibanserin acts on serotonin (a chemical produced in the brain). Serotonin is believed to play an important role in the regulation of mood, sleep, sexuality, and appetite.

This clinical trial is being conducted at: Mount Carmel East  
The Physician's Office

For more information about this clinical trial call: (614) 861-2707

